




## Synapses, Systems and Simulations



Nicolas Le Novère Babraham Institute, n.lenovere@gmail.com





Duration: decades

- Amyloid-β accumulates in the brain
- Tau hyperphosphorylation gradually leads to neuronal loss
- · Pathology does not yet noticeably affect cognition
- · Biomarkers and genetic profile can indicate the risk of disease progression and reveal underlying AD

Subtypes:


- Progressive MCI
- MCI caused by AD
- · Amnestic syndrome of the hippocampal type
- · Deficits in memory and/or other cognitive domains noticeable to the person affected and/or others, but not severe enough to interfere with activities of daily living
- · Biomarkers can determine the aetiological diagnosis

Duration: ~7 years; followed by total loss of independent function

- · Notable loss of intellectual ability affecting memory and at least one other cognitive domain
- The impairment interferes with activities of daily living

Hampel and Lista (2016) Nature Reviews | Neurology





interfere with activities of

 Biomarkers can determine the aetiological diagnosis

daily living

Hampel and Lista (2016) Nature Reviews | Neurology




of disease progression and

reveal underlying AD

"Rather than neuronal loss, it is now widely accepted that synaptic impairments underlie the decrease performance"


Federico Grillo - 2016






Fleming and England (2010) Nat Chem Biol






Fleming and England (2010) Nat Chem Biol






Fleming and England (2010) Nat Chem Biol







lasts for weeks


Fleming and England (2010) Nat Chem Biol



#### The calcium theory of synaptic plasticity L-type **NMDAR AMPAR** VOC **SCENARIO** ER Lisman (1989) PNAS, 86: 9574-8 Calmodulin low high calcium calcium Calcineurin (PP2B, PP3)d CaMKII **LONG TERM LONG TERM**



**DEPRESSION** 









Calcium
















#### Calmodulin carries 4 Ca<sup>2+</sup> binding domains





















































#### Targets as allosteric effectors: the evidence




Peersen et al. (1997)



## Bindings of calcium and protein targets







Calcium

Conf. transition

Calcium binding

Target binding

$$c = \frac{K_R}{K_T}$$

$$e = \frac{K_R^t}{K_T^t}$$



## Targets as allosteric effectors: the theory

$$\bar{Y}_{\text{Ca}^{2+}} = \frac{\alpha(1+\alpha)^{n-1} + L'c\alpha(1+c\alpha)^{n-1}}{(1+\alpha)^n + L'(1+c\alpha)^n}$$

$$L = \frac{[T_0]}{[R_0]}$$

$$L' = L\left(\frac{1+\gamma e}{1+\gamma}\right)^n$$

$$c = \frac{K^R}{K^T}$$


$$e = \frac{K_t^R g}{K_t^T g}$$

$$\alpha = \frac{[Ca^{2+}]}{[K^R]}$$

$$\gamma = \frac{[Target]}{[K_t^R g]}$$




## Different targets stabilise lobes in different states



Lai M, Brun D, Edelstein SJ, Le Novère N (2015) Lai M, Edelstein SJ, Le Novère N (in preparation)




#### Hemiconcerted model of calmodulin






## Calcium binding to lobes and whole CaM (exp)





## Calcium binding to lobes and whole CaM (sim)





[CaM]

#### Parametrisation using accurate measurements

- Ca<sup>2+</sup> binding in presence of targets: none, skMLCK, PhK5, CaATPase
- Ca<sup>2+</sup> dissociation constants for complete calmodulin and N and C term mutants



$$C=3.96\ 10^{-3}$$

Affinity of Ca<sup>2+</sup> for "open state" 250 times higher than for "closed state"

$$K_A^R = 8.32 \ 10^{-6}$$
  
 $K_B^R = 1.66 \ 10^{-8}$   
 $K_C^R = 1.74 \ 10^{-5}$   
 $K_D^R = 1.45 \ 10^{-8}$ 

2 high, 2 low, as expected



## Activity of unsaturated calmodulin (state function)


Fractional activity depends on the number of calcium ions bound

$$\frac{R_2}{T_2} = \frac{1}{L \cdot c^2}$$

- $R_0/T_0 = 1/20000 (1/L)$
- $R_1/T_1 = 1/170$
- $R_2/T_2 = 0.69$  half-saturation  $\approx$  equi-probability
- $R_{4}/T_{4} = 10000$




## Effect of an R-stabilizing target on Ca<sup>2+</sup> affinity






## Binding to target increases the affinity for Ca<sup>2+</sup>






# Targets stabilises Ca<sup>2+</sup> binding into the physiological range






## Calmodulin, its ligand and its targets






## **Bidirectional synaptic plasticity**






### **Calcineurin stabilises CaM R** → no deactivation







# Neurogranin binds to apo-CaM, decrease affinity for [Ca<sup>2+</sup>] and increase dissociation rate




Kumar *et al* (2013)




### Effect of R and T stabilising targets on CaM affinity













### No large effect of Ng on [Ca2+free]






## Ng affects the distribution of CaM









# Wait a minute! Signal transduction is not at equilibrium!

AMPAR post-synaptic potential: 5 ms

Calcium spike: 50 ms

Half saturation calmodulin (kon=1.5e6, koff=100): 5 ms

Relaxation between calmodulin states: 1 ms


autophosphorylation of CaMKII (kon=6): 100 ms











#### Calcium/calmodulin kinase II


catalytic site **Auto-inhibitory** domain T306P blocks CaM binding CaM binding site T286P causes constitutive activity

Calmodulin trapping is an apparent increase of affinity of CaMKII for CaM when T286 is phosphorylated

Stefan MI, Marshall D, Le Novère N (2012)




#### Calcium/calmodulin kinase II




Dodecamer:


Trans-phosphorylation of T286 by neighbouring subunits Cis-phosphorylation of T306


Most quantitative measurements made on monomers ...













Bradshaw JM, Kubota Y, Meyer T, Schulman H (2003). *PNAS* 100: 10512–10517.






Black DJ, Selfridge JE, Persechini A (2007). *Biochemistry* 46: 13415–13424.

#### Validation of calciumactivation of CaN

Quintana AR, Wang D, Forbes JE, Waxham MN (2005). *BBRC* 334: 674–680.







## **Calmodulin** activation

CaM without targets. Low frequencies do not activate calmodulin (binding events without conformational changes)







0.1 -60

120

60


time (s)

frequency (Hz)

→ bistability

Positive feedback loop

## Temporal activation of CaMKII and CaN






6

5

### **Bidirectional plasticity**



60

Constant catalytic rates of active enzyme

→ quantity of catalysed reaction events prop to integral of the activation curve




### **Bidirectional plasticity**




→ quantity of catalysed reaction events prop to integral of the activation curve

Bienestock-Cooper-Munro (BCM) curve: difference of active areas\*catalytic activities









### **Bidirectional plasticity**


Constant catalytic rates of active enzyme

→ quantity of catalysed reaction events prop to integral of the activation curve

Bienestock-Cooper-Munro (BCM) curve: difference of active areas\*catalytic activities











#### Effect of calcium duration and amount




Prolonged or intense signals decrease  $\Theta$ m: It is not an intrinsic property of the synapse





### Effect of intrinsic system perturbations










Scilab









Lu Li



Denis Brun

### wellcome trust





Stuart Edelstein



Massimo Lai











